

RESEARCH

Open Access

A novel whole-cell lysate kinase assay identifies substrates of the p38 MAPK in differentiating myoblasts

James DR Knight^{1,2,9}, Ruijun Tian^{3,4,9}, Robin EC Lee^{1,2,10}, Fangjun Wang^{3,5}, Ariane Beauvais¹, Hanfa Zou⁵, Lynn A Megeney^{1,2,6}, Anne-Claude Gingras^{7,8}, Tony Pawson^{7,8}, Daniel Figey^{3,4} and Rashmi Kothary^{1,2,6*}

Abstract

Background: The p38 α mitogen-activated protein kinase (MAPK) is a critical mediator of myoblast differentiation, and does so in part through the phosphorylation and regulation of several transcription factors and chromatin remodelling proteins. However, whether p38 α is involved in processes other than gene regulation during myogenesis is currently unknown, and why other p38 isoforms cannot compensate for its loss is unclear.

Methods: To further characterise the involvement of p38 α during myoblast differentiation, we developed and applied a simple technique for identifying relevant *in vivo* kinase substrates and their phosphorylation sites. In addition to identifying substrates for one kinase, the technique can be used *in vitro* to compare multiple kinases in the same experiment, and we made use of this to study the substrate specificities of the p38 α and β isoforms.

Results: Applying the technique to p38 α resulted in the identification of seven *in vivo* phosphorylation sites on six proteins, four of which are cytoplasmic, in lysate derived from differentiating myoblasts. An *in vitro* comparison with p38 β revealed that substrate specificity does not discriminate these two isoforms, but rather that their distinguishing characteristic appears to be cellular localisation.

Conclusion: Our results suggest p38 α has a novel cytoplasmic role during myogenesis and that its unique cellular localisation may be why p38 β and other isoforms cannot compensate for its absence. The substrate-finding approach presented here also provides a necessary tool for studying the hundreds of protein kinases that exist and for uncovering the deeper mechanisms of phosphorylation-dependent cell signalling.

Keywords: differentiation, FSBA, kinase assay, mitogen-activated protein kinase, myoblast, p38, phosphorylation, quantitative MS

Background

Protein kinases are well-known regulators of cell signalling and cellular behaviour that execute their function through the covalent attachment of an ATP-derived phosphate to protein substrates. To understand the function of any protein kinase on a large and cell-wide scale first requires the development of a substrate screening technique that allows for the proteins phosphorylated by a kinase of interest to be comprehensively identified, ideally in a single experiment. Although

substrate-finding techniques exist, they are hindered by problems that prevent them from being easily or readily employed [1-4] and are generally limited to providing *in vitro* substrate identifications that may or may not be relevant *in vivo*. *In vivo* approaches currently available, such as that employed by Holt *et al.* [5], can associate a kinase with *in vivo* phosphorylation events, but direct phosphorylation cannot be inferred without additional experimentation. A simple technique that can identify direct *in vivo* substrates is an obvious need for the field.

The mitogen-activated protein kinase p38 α is involved in several cellular processes, but its critical role during differentiation, and particularly the differentiation of myoblasts, has been a major focus. At the initiation of

* Correspondence: rkothary@ohri.ca

¹Regenerative Medicine Program, Ottawa Hospital Research Institute, 501 Smyth Road, Ottawa, ON, K1H 8L6, Canada

Full list of author information is available at the end of the article

myoblast differentiation, p38 α is known to phosphorylate several transcription factors and chromatin remodelling proteins, thereby inducing the expression of a myogenic gene program [6]. Although much is known about p38 α 's role in this process, it is likely very partial, and whether p38 α plays an important role in other processes during myoblast differentiation, such as cell fusion or sarcomere formation, is unknown. At the same time, there are questions regarding the other p38 isoforms and their role, or lack thereof, in myogenesis. p38 β is also expressed in myoblasts and is activated in the same manner as p38 α , but despite having a kinase domain 75% identical to that of p38 α (72% sequence identity overall), p38 β is unable to compensate for the loss of p38 α , even when overexpressed [7-9]. The obvious and suspected explanation is that there are critical myogenic phosphorylations specific to the α isoform, but these have yet to be discovered and whether this assumption is correct is unknown.

Here we describe a simple approach for substrate finding that can be used to identify *in vitro* and *in vivo* substrates. The technique begins with treatment of cell lysate to inactivate endogenous kinases, followed by an *in vitro* assay using an exogenous kinase of interest, and concludes with quantitative mass spectrometry (MS) to identify phosphorylation sites specific to the added kinase. By using lysate derived from vehicle- or inhibitor-treated cells, this *in vitro* approach can be simultaneously coupled with biologically relevant information to identify direct substrates regulated by the kinase of interest *in vivo*. Applying this technique to p38 α with lysate from differentiating myoblasts resulted in the identification of several new *in vivo* substrates that suggest novel functions for p38 α during myogenesis. We did not identify a single phosphorylation specific to the p38 α isoform compared with p38 β , at least in terms of *in vitro* substrate specificity, but we did see a clear difference in cellular localisation during myoblast differentiation. This leads us to propose that although the kinase domains of p38 α and β likely have the same capacity to phosphorylate substrates, there are major differences in actual substrate specificity in an *in vivo* context.

Methods

Cell culture

C2C12 cells were grown in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% (vol/vol) foetal bovine serum with 100 U/ml penicillin, 100 μ g/ml streptomycin and 250 ng/ml amphotericin B. To induce differentiation, cells were grown to 85-90% confluence, and the medium was changed to DMEM with 2% horse serum supplemented with penicillin, streptomycin and amphotericin B as described above. FLAG-tagged p38 α

and p38 β were acquired from Addgene (Cambridge, MA, USA) [10]. Constructs were transfected into C2C12 myoblasts with Lipofectamine 2000 reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. To inhibit p38 activity, SB 202190 (Promega, Madison, WI, USA) solubilised in dimethyl sulphoxide (DMSO) or DMSO as control was added to differentiation media at 48 hours following the induction of differentiation, and medium with inhibitor was changed daily.

Immunofluorescence

Cells were fixed with 4% formaldehyde and stained with the following antibodies: Flag M2 (1:1,000 dilution; Sigma-Aldrich, St Louis, MO, USA), myosin heavy chain (MyHC) (1:20 dilution; Developmental Studies Hybridoma Bank, Department of Biology, University of Iowa, Iowa City, IA, USA), Alexa Fluor 488 goat anti-mouse antibody (1:1,000 dilution; Molecular Probes/Invitrogen) and Alexa Fluor 555 goat anti-mouse antibody (1:1,000 dilution; Molecular Probes/Invitrogen). The differentiation index was calculated as the number of MyHC-positive nuclei divided by the total number of nuclei. The fusion index was quantified as the number of nuclei per MyHC-positive cell. Five fields of view at $\times 20$ magnification were counted and averaged per replicate, with a total of three replicate experiments.

Statistical analysis

Statistical analyses were performed using StatPlus software (AnalystSoft Inc; <http://www.analystsoft.com/en/products/statplus/>). The data shown are means with SD, and Student's *t*-tests were performed to determine significance for the differentiation and fusion indices.

Western blot analysis

For Western blot analysis, cells were lysed in radioimmunoprecipitation assay (RIPA) buffer (50 mM Tris-HCl, pH 7.5, 1% Nonidet P-40, 0.1% SDS, 150 mM NaCl, 1 mM ethylenediaminetetraacetic acid (EDTA), 50 mM NaF, 200 μ M Na₃VO₄, 1 mM phenylmethanesulphonylfluoride (PMSF), 10 μ g/ml aprotinin, 10 μ g/ml leupeptin and 10 μ g/ml pepstatin), and 5 \times Laemmli buffer (300 mM Tris, pH 6.8, 0.01% bromophenol blue, 10% SDS, 50% glycerol and 5% β -mercaptoethanol) was added to 25 μ g of protein per sample to a final concentration of 1 \times for SDS-PAGE. For fractionation, cells were lysed in hypotonic buffer (10 mM 4-(2-hydroxyethyl)-1-piperazineethanesulphonic acid (HEPES), pH 7.5, 1.5 mM MgCl₂, 10 mM KCl, 0.5 mM dithiothreitol (DTT), 50 mM NaF, 200 μ M Na₃VO₄, 1 mM PMSF, 10 μ g/ml aprotinin, 10 μ g/ml leupeptin and 10 μ g/ml pepstatin) and left on ice for 10 minutes. Cells were then passed through a 25-gauge needle three times and

centrifuged at 500 g to pellet nuclei and unlysed cells. The supernatant was collected as whole cytoplasm. For further fractionation of the cytoplasm, the supernatant was centrifuged again at 5,000 g to pellet mitochondria and membrane fractions. The supernatant was then collected and centrifuged at 100,000 g to pellet any remaining cell particles, and the resulting supernatant was collected as cytosol. RIPA buffer was added to all fractions to a final concentration of 1× for complete lysis. Recombinant phosphorylated p38 α (Millipore, Billerica, MA, USA) was dephosphorylated using λ -protein phosphatase (New England Biolabs, Inc, Ipswich, MA, USA) by adding 800 U of phosphatase to 200 ng of p38 α diluted in λ -phosphatase buffer, and the sample was assayed at 30°C for 1 hour. Antibodies used for blotting were as follows: α -actinin (1:125 dilution; Abcam, Cambridge, MA, USA), COX IV (1:1,000 dilution; Abcam), GRP78 (1:500 dilution; Cell Signaling Technology, Danvers, MA, USA), lamin A/C (1:500 dilution; Abcam), MyHC (1:100 dilution; Developmental Studies Hybridoma Bank), MyoD (1:1,000 dilution; Santa Cruz Biotechnology, Santa Cruz, CA, USA), myogenin (1:100 dilution; Developmental Studies Hybridoma Bank), neural cell adhesion molecule (1:200 dilution; Abcam), p38 α (1:500 dilution; Cell Signaling Technology), phospho-p38 (1:500 dilution; Abcam) and β -tubulin (1:1,000 dilution; Developmental Studies Hybridoma Bank). An Alpha Innotech HD2 imaging system (R&D Systems, Minneapolis, MN, USA) was used to quantify phospho-p38 and tubulin expression.

FSBA treatment and substrate labelling

Cells were lysed in Nonidet P-40 buffer (50 mM Tris-HCl, pH 7.8, 150 mM NaCl, 1% (vol/vol) Nonidet P-40, 1 mM PMSE, 10 μ g/ml aprotinin, 10 μ g/ml leupeptin and 10 μ g/ml pepstatin). Lysate was treated at a concentration of 2 mg/ml with 20 mM 5'-4-fluorosulphonylbenzoyl adenosine (FSBA) solubilised in DMSO and placed at 30°C for 1 hour. The sample was then diluted down to 1:5 with Nonidet P-40 buffer minus protease inhibitors and desalted using Millipore Amicon ultrafiltration columns with a 10 kDa molecular weight cutoff. Following concentration, the sample was diluted to 4 mg/ml with Nonidet P-40 buffer and diluted 1:2 with 2× kinase assay buffer (40 mM 3-morpholinopropane-1-sulphonic acid (MOPS), pH 7.2, 50 mM β -glycerophosphate, 10 mM ethylene glycol tetraacetic acid (EGTA), 2 mM Na₃VO₄, 2 mM DTT, 50 mM MgCl₂, 400 μ M cold ATP and 5 μ Ci [γ -³²P]ATP). Recombinant p38 α or p38 β (Millipore) was added to a final concentration of 0.5% (wt/wt) total protein. Control and kinase-added samples were assayed at 30°C for 1.5 hours. For one-dimensional SDS-PAGE, 5× Laemmli buffer was added following the assay to 1×, and the

sample was electrophoresed. For two-dimensional electrophoresis, 17-cm ReadyStrip immobilised pH gradient (IPG) strips (Bio-Rad Laboratories, Hercules, CA, USA) were directly rehydrated with labelled lysate diluted in rehydration buffer (7 M urea, 2 M thiourea, 4% 3-[(3-cholamidopropyl)dimethylammonio]-1-propanesulphonate (CHAPS) and 1% DTT) following the manufacturer's directions. Isoelectric focusing was performed on a PROTEAN IEF Cell (Bio-Rad Laboratories) under the following conditions: 200 V for 1 hour, 500 V for 1 hour, 5,000 V ramp for 5 hours and 5,000 V for 80,000 VH. IPG strips were then equilibrated following the manufacturer's instructions and overlaid onto a 12% SDS-PAGE gel. Following electrophoresis, gels were dried and imaged. For one-dimensional electrophoresis, 100 μ g of lysate was used per reaction. For two-dimensional electrophoresis, 300 μ g of lysate was used.

For *in vitro* substrate identification, assays were performed as described above with the following modifications. A quantity of 1.5 mg of lysate was treated with FSBA, the sample was desalted and 2× kinase assay buffer was added (40 mM MOPS, pH 7.2, 50 mM β -glycerophosphate, 10 mM EGTA, 2 mM Na₃VO₄, 2 mM DTT, 50 mM MgCl₂ and 2 mM cold ATP). The sample was then split into three 500- μ g aliquots, and 5 μ g of heat-inactivated p38 α was added to the control, 5 μ g of active p38 α was added to the second aliquot and 5 μ g of active p38 β was added to the third aliquot. The samples were then assayed for 3 hours at 30°C.

For *in vivo* substrate identification, assays were performed as above with the following modifications. At 48 hours of differentiation, cells were treated with 5 μ M of SB 202190 or an equivalent amount of DMSO as vehicle. Twenty-four hours later the cells were lysed in Nonidet P-40 buffer. Lysate (1 mg) from DMSO-treated cells or 2× 1 mg of lysate from SB 202190-treated cells was inhibited with FSBA, the samples were desalted and 2× kinase assay buffer was added to each. A quantity of 5 μ g of active p38 α was added to one of the lysate aliquots from SB-treated cells. The samples were then assayed for 3 hours at 30°C.

Dimethyl labelling

After assaying the samples, they were precipitated by methanol chloroform, then redissolved in 200 μ l of 8 M urea and 50 mM Tris-HCl, pH 8.1, with sonication. The samples were then reduced with 20 mM DTT for 1 hour at 60°C and alkylated by 100 mM iodoacetamide for 30 minutes at room temperature in the dark. Subsequently, the samples were diluted to 2 M urea with 50 mM Tris-HCl, pH 8.1, and digested with trypsin at a protein-to-trypsin ratio of 50:1 (wt/wt) for 16 hours at 37°C. Next, the digested samples were acidified to pH 2 using 10% (vol/vol) formic acid. Dimethyl labelling of

the samples was performed as reported previously [11] and is described briefly as follows. The acidified peptides were loaded onto C18 solid phase extraction (SPE) columns (50 mg of packing material). After brief washing with 50 mM sodium phosphate buffer, pH 7.5, 3 ml of light, intermediate and heavy labelling reagents were loaded onto C18 SPE columns trapped with control, p38 α - and p38 β -labelled samples, respectively. After being washed with 0.1% (vol/vol) formic acid, the labelled samples were eluted with 80% acetonitrile (ACN) (vol/vol) and 0.1% (vol/vol) formic acid, then dried by vacuum centrifugation.

Phosphopeptide enrichment

Phosphopeptide enrichment by TiO₂ was carried out as reported previously [12] with modifications. The dried samples were redissolved with 65% ACN/2% trifluoroacetic acid (TFA)/saturated glutamic acid and combined. TiO₂ beads suspended in 65% ACN/2% TFA/saturated glutamic acid were added into the above samples with a peptide to TiO₂ bead ratio of 1:4 (wt/wt). After being nutated for 40 minutes, the TiO₂ beads were recovered by centrifugation and washed thoroughly with 65% ACN/2% TFA. Finally, the enriched phosphopeptides were eluted with 10% (vol/vol) NH₃·H₂O and dried by vacuum centrifugation.

Online liquid chromatography tandem mass spectrometry analysis

Online liquid chromatography tandem mass spectrometry (LC-MS/MS) analysis was performed as reported previously [13,14] with modifications. The dried sample was redissolved with 0.1% formic acid and loaded onto a biphasic trap column (200 μ m ID \times 10 cm; 5-cm reversed phase column packed with ReproSil-Pur C18 resin (5 μ m at 200 Å ; Dr.Maisch GmbH, Ammerbuch-Entringen, Germany) and a 5-cm monolith strong cation exchange (SCX) column). The trapped phosphopeptides were eluted from the trap column onto a C18 tip column (75 μ m ID \times 20 cm, 3 μ m at 200 Å ; Dr.Maisch GmbH) by a series of salt washes at increasing concentrations (0, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 70, 80, 90, 100, 150, 200 and 1,000 mM). Each fraction was then separated by reversed phase-based gradient elution and detected using an LTQ Orbitrap XL Fourier transform mass spectrometer (Thermo Scientific, Waltham, MA, USA). The reversed phase gradient was set as follows: 0% to 5% ACN for 2 minutes, 5% to 30% ACN for 90 minutes and 30% to 80% ACN for 5 minutes. After flushing with 80% ACN for 10 minutes, the column was equilibrated with 0.1% formic acid aqueous solution for 13 minutes. The LTQ Orbitrap XL Fourier transform mass spectrometer was operated in positive ionization mode. A voltage of 1.8 kV was applied. MS

and MS/MS spectra were acquired in a data-dependent mode, and one full MS scan was followed by ten MS/MS scans. The resolution was set at 60,000 at m/z 400 after accumulation to a target value of 500,000.

Protein identification and quantification

All MS/MS spectra in one acquired raw file were converted to a single *.mgf file using DTASuperCharge version 2.0a7 (Matrix Science, Boston, MA, USA). The *.mgf file was queried against the International Protein Index mouse database version 3.52 (EMBL-EBI; <http://www.ebi.ac.uk/IPI/IPIhelp.html>) using Mascot version 2.1 (Matrix Science). To evaluate the false discovery rate (FDR), reversed sequences were appended to the database. Cysteine residues were searched as a static modification of +57.0215 Da; methionine residues were searched with a variable modification of +15.9949 Da; and serine, threonine and tyrosine residues were searched with a variable modification of +79.9663 Da. Light, intermediate and heavy dimethylation of peptide amino termini and lysine residues were set as variable modifications of +28.0313 Da, +32.0564 Da and +36.0757 Da, respectively. Peptides were queried using full tryptic cleavage constraints with up to two missed cleavage sites. The mass tolerances were 7 ppm for parent masses and 0.5 Da for fragment masses. Phosphopeptides with a Mascot score \geq 30 (rank 1, $P \leq$ 0.05, bold red required) were selected and quantified (FDR < 0.01). Phosphorylation site localisation and phosphopeptide quantification were performed using a dimethyl-adapted version of MSQuant version 2.0a81. For each peptide, the putative site of phosphorylation yielding the highest posttranslational modification (PTM) score was accepted (PTM score > 13 required, as described previously [15]). Peptide ratios were obtained by calculating the extracted ion chromatograms of the light, medium and heavy forms of the peptide using the monoisotopic peaks only, and protein ratios were calculated from the average of all quantified peptides. All MSQuant outputs of the same online multidimensional separation were then imported into StatQuant version 1.2.2, and the quantified phosphopeptides were sorted together and exported.

Substrate validation

1 μ g recombinant HSP27 (Enzo Life Sciences, Farmingdale, NY, USA), recombinant Rps27 (Abnova, Taipei City, Taiwan) or recombinant Smad9 (Abnova) were diluted in 30 μ l of 1 \times kinase assay buffer (20 mM MOPS, pH 7.2, 25 mM β -glycerophosphate, 5 mM EGTA, 1 mM Na₃VO₄, 1 mM DTT, 25 mM MgCl₂, 200 μ M cold ATP and 2.5 μ Ci [γ -³²P]ATP). A quantity of 500 ng of p38 α or p38 β was then added, and the samples were assayed at 30°C for 1 hour. Laemmli buffer at

5× concentration was added to 1× to terminate the assays, then the samples were electrophoresed and the gels were dried and imaged. Peptide assays were performed similarly with 5 µg of the following peptides synthesised by Biomatik (Cambridge, ON, Canada): HSP27 S180-APLPKAVTQSAEITIPVTF, HSP27 A180-APLPKAVTQAAEITIPVTF, Rps27 S27-KHKKKRLVQ SPNSYFMDVK, Rps27 A27-KHKKKRLVQAPNSYF MDVK, Smad9 T136-NPYHYQRVETPVLPPVLVP and Smad9 A136-NPYHYQRVEAPVLPPVLVP. The frequency logo was generated using WebLogo [16].

Results

FSBA inhibits endogenous protein kinases

A substrate-finding approach that works with cell lysate must overcome the obstacle of endogenous protein kinase activity. Lysate contains tens if not hundreds of active kinases, making it difficult to attribute individual phosphorylations that occur during a lysate-based assay to a particular kinase. 5'-4-fluorosulphonylbenzoyl adenosine (FSBA) offers a simple solution. FSBA is an ATP analogue that inhibits protein kinases by occupying the ATP binding site and covalently attaching to an invariant lysine [17-20], the fully functionally conserved and so-called catalytic lysine [21]. As FSBA irreversibly occupies the ATP binding site, a bound kinase will permanently lose activity. Treatment of whole-cell C2C12 myoblast lysate with this compound can completely eliminate the endogenous kinase signal present (Figure 1A).

Kinase-specific substrate labelling

Cell lysate treated with FSBA can be desalted to remove any unbound inhibitor and a pool of protein with no inherent kinase activity is generated. A kinase of interest can then be added with a kinase assay buffer, and any labelling that subsequently occurs is due to the added kinase as opposed to an endogenous one. To specifically label substrates of p38α, a kinase assay buffer containing [γ -³²P]ATP was added to FSBA-treated C2C12 lysate, along with recombinant p38α, and the sample was assayed at 30°C. Substrates labelled by p38α appeared as bands following one-dimensional gel electrophoresis or as spots in two-dimensional gel electrophoresis with no contaminating signal from endogenous kinases (Figure 1B and Additional file 1 Figure S1). Although this type of approach is excellent for visualising phosphorylation, it is very difficult to identify phosphorylated proteins through spot-picking and MS. We therefore sought an alternative gel- and radioactive-free approach for identifying phosphorylated proteins.




Figure 1 FSBA is a pan-kinase inhibitor that allows for kinase-specific substrate labelling of cell lysate. **(A)** C2C12 cell lysate was treated with either nothing (lane 1), dimethyl sulphoxide (DMSO) or 5'-4-fluorosulphonylbenzoyl adenosine (FSBA) solubilised in DMSO. Subsequently, samples were desalted, kinase assay buffer containing ³²P- γ -ATP was added and the samples were assayed for 1.5 hours. Treatment with FSBA abolished the labelling of endogenous protein kinase substrates. **(B)** After pretreatment of C2C12 cell lysate with FSBA, purified p38α was added with a kinase assay buffer to specifically label its substrates and visualised using one-dimensional SDS-PAGE or two-dimensional gel electrophoresis (Additional file 1 Figure S1).





Figure 2 Methodology used to identify *in vitro* kinase substrates using FSBA and quantitative mass spectrometry. The differential tagging of peptides with dimethyl labels allows for the relative quantification and identification of phosphopeptides simultaneously in a high-mass accuracy mass spectrometer. An example/theoretical substrate identification for the added kinase (green) is shown and is based on the higher intensity of the phosphopeptide in MS.

Quantitative MS coupled with a phosphopeptide enrichment to identify substrates


The approach we devised to identify substrates is outlined in Figure 2. Cell lysate is treated with FSBA and desalted as described in the previous section. This is followed by the addition of a nonradioactive kinase assay buffer, and the sample is split in two, to one of which is added active kinase (kinase-added). After assaying at 30°C, the two samples are digested, and peptides from each sample are differentially tagged using isotopomeric

dimethyl labels. As a final step before MS, the samples are combined and an enrichment for phosphopeptides is performed using either TiO_2 for serine/threonine phosphorylation or a phosphotyrosine antibody for tyrosine phosphorylation. Phosphopeptides are then identified by performing LC-MS/MS, and their relative abundance between samples quantified by the differential dimethyl labelling of peptides. Phosphopeptides that are more abundant in the kinase-added sample are from proteins labelled by the added kinase during the *in vitro* assay, and by this means substrates can be identified. This approach results not only in substrate identification but also in the identification of the site of phosphorylation. Because up to three samples can be compared using dimethyl labelling, the phosphorylation profile of two kinases (plus a control) can be compared in a single experiment.

We began our initial *in vitro* substrate-finding procedure by treating 1.5 mg of C2C12 lysate with FSBA. For the assay, the sample was split into three equal parts, with active p38 α and β added to the second and third samples, respectively. The rationale behind comparing p38 α with p38 β was to identify specific p38 α phosphorylations that might explain why p38 β cannot compensate for the loss of p38 α in differentiating myoblasts. In total, 387 unique serine/threonine phosphopeptides were identified (Additional file 1 Table S1). A histogram of their relative abundance ratios (p38 α /control) is shown in Additional file 1 Figure S2. A threefold increase in the abundance ratio was selected as a cutoff for high-confidence *in vitro* substrates as this ratio is beyond the range of inherent variability, and 158 phosphorylation sites from 94 different proteins showed at least a threefold increase in the p38 α -assayed sample relative to the control. The list of p38 α phosphorylation sites is presented in Additional file 1 Table S2. We identified five previously known substrates (caldesmon, histone H2B, *Psmc1*, SAKS1 and Smad3) and eighty-nine that were previously unknown. Three of these previously unknown targets were validated to determine if the technique was discovering true *in vitro* p38 α substrates. Recombinant forms of these three proteins could be phosphorylated *in vitro* by p38 α (Figure 3A), and a validation of peptides confirmed the site of phosphorylation (Figure 3B). After aligning substrates on their phosphorylation sites, we found that a consensus phosphorylation motif was present in many substrates, although it was not an absolute requirement (Figure 3C). The motif contains a proline immediately downstream from the target serine or threonine and an aliphatic residue two residues upstream. This motif is in agreement with that previously described for p38 [22], providing further support for our substrate-finding approach.



Substrate specificity does not distinguish p38 α and p38 β
 We first assayed p38 α and p38 β on cell lysate using the radioactive approach and were surprised to see no obvious differences in the substrate banding patterns produced by the two isoforms (Figure 4A). Consistent with this observation, of the 158 p38 α phosphorylations identified using the quantitative approach, none appeared to be specific to this isoform. The 158 quantitative values for the p38 α phosphorylations are plotted from highest to lowest in Figure 4B (blue line for p38 α). The red line represents the corresponding p38 β values. Although p38 α and p38 β showed apparent differences in preference, they had very similar overall profiles. There were only four phosphorylations that fell above the threefold cutoff for p38 α , whereas for p38 β they were well below. However, the values were still positive for p38 β , suggesting that these could be real phosphorylations but there is simply less confidence in them. To determine if these might represent specific phosphorylations, we performed *in vitro* assays with either p38 α or p38 β using purified substrate for one candidate. The purified substrate was the 40S ribosomal protein S27 (Rps27), and both p38 isoforms were able to phosphorylate this protein at the same site (Figure 4C and 4D).



Therefore, none of the 158 phosphorylations found for p38 α appear to be specific to this isoform.

If substrate specificity does not distinguish the two p38 isoforms, and yet p38 β is unable to compensate for the loss of p38 α , there must be an alternative characteristic that discriminates them. An obvious possibility is cellular localisation. If p38 α and p38 β localise differently within the cell, then p38 β would simply be unable to fulfil p38 α 's role, even though it might have the catalytic potential to do so. To study this, we overexpressed FLAG-tagged p38 α and p38 β in C2C12 cells and assessed their localisation during differentiation (Figure 4E). Whereas p38 α has a ubiquitous localisation pattern, p38 β is found only at the periphery of the cell. p38 α therefore has access to a substrate pool that p38 β does not, highlighting a major reason why p38 β cannot compensate for the loss of p38 α .

Identification of novel *in vivo* p38 α substrates

Although the approach we have outlined can identify *in vitro* kinase substrates, such *in vitro* approaches will also turn up irrelevant substrates that are phosphorylated only because of the absence of the appropriate cellular context. For example, we identified just over 400 phosphorylation sites in the screen described above, 158 of which are *in vitro* p38 targets. If these numbers are representative, it suggests that p38 could be responsible for approximately 40% of all cellular phosphorylations in differentiating myoblasts, a number that seems unlikely, given that there are about 500 mouse and human kinases [23].

To overcome the drawbacks associated with *in vitro* substrate identification, we modified the approach to allow for the identification of relevant, direct *in vivo* substrates (outlined in Figure 5). For our purposes, differentiating C2C12 cells were treated either with the p38 inhibitor SB 202190 or with DMSO as vehicle prior to lysis. SB inhibitor or vehicle treatment began at 48 hours of differentiation, and the cells were lysed 24 hours later. The 48-hour time point was chosen because of our interest in identifying novel functions for p38 α during the middle stages of myoblast differentiation. Myogenic gene activation occurs within the first 48 hours of differentiation [24] and is followed by cell fusion, sarcomere formation and other processes. If differentiating C2C12 myoblasts are treated with a p38 inhibitor at 48 hours, there is a reduction in cell fusion and overall differentiation (Additional file 1 Figure S3), indicating that there is a requirement for long-term p38 activity for efficient myogenesis beyond the initial stage of myogenic gene activation, possibly both for maintaining such gene activation and for fusion-related processes. Following cell lysis, a 1-mg aliquot was taken from the lysate of vehicle-treated cells and two 1-mg




Figure 5 Methodology used to identify *in vivo* kinase substrates using FSBA and quantitative MS. *In vivo* substrates are identified based on the higher phosphopeptide intensity in MS for both the vehicle-treated (blue) and kinase-added (green) samples. A higher phosphopeptide abundance in MS for the vehicle-treated (blue) sample relative to the inhibitor-treated (red) sample indicates a phosphorylation site regulated by the kinase of interest *in vivo*. A higher phosphopeptide abundance for the kinase-added (green) sample relative to the inhibitor-treated (red) sample indicates a phosphorylation site directly targeted by the kinase *in vitro*.

aliquots were taken from the lysate of inhibitor-treated cells. All three aliquots were treated with FSBA and desalted to remove unbound FSBA and SB inhibitor, and a kinase assay buffer was added to each. Active p38 α was added to one of the inhibitor-treated aliquots, and the samples were assayed at 30°C. Following the assay, the samples were digested, dimethyl-labelled and phosphopeptides enriched as described above. What we sought to identify were phosphorylation sites that decreased on inhibitor treatment and could be elevated back up by direct *in vitro* p38 α labelling of SB inhibitor-treated lysate. These would be direct phosphorylation sites regulated by p38 α *in vivo*.

By using this approach, we identified a total of 717 phosphorylation sites (Additional file 1 Table S3), 73 of


Table 1 Newly discovered *in vivo* p38 α substrates

Genes	Protein names	Sites
<i>Ahnak</i>	Neuroblast differentiation-associated protein <i>AHNAK</i>	S4906
<i>Iws1</i>	Protein IWS1 homolog	S183, S185
<i>Grp78</i>	78 kDa glucose-regulated protein	T649
<i>Pgrmc</i>	Membrane-associated progesterone receptor component 1	S181
<i>Prdx6</i>	Peroxiredoxin 6	T93
<i>Ranbp2</i>	E3 SUMO ^a protein ligase RanBP2	S2505

^aSUMO, small ubiquitin-like modifier.

which showed a threefold decrease following SB inhibitor treatment, and seven of these sites were direct p38 α phosphorylation sites (that is, showed at least a threefold increase in the p38 α -labelled sample relative to the lysate from inhibitor-treated cells). In contrast to the pure *in vitro* approach, which suggested that approximately 40% of all cellular phosphorylations might be p38 target sites, the numbers derived from the *in vivo* study suggest that the actual level is closer to about 1%. The seven *in vivo* sites (on six substrates) that we identified are listed in Table 1.

The substrate *Iws1* is a largely nuclear protein involved in several processes, such as chromatin remodelling and mRNA export [25,26], functions with which p38 has previously been associated. *Ranbp2* localises predominantly to the nuclear pore complex and is critical for myotube formation [27]. The other four substrates are cytoplasmic proteins without known or suggested nuclear functions (*Ahnak*, *Grp78*, *Pgrmc* and *Prdx6*). On the basis of fractionations and Western blot analysis, we found that p38 α was indeed present in the cytoplasm during differentiation (Figure 6A), which is in agreement with the immunofluorescence staining of FLAG-tagged p38 α . The levels of phosphorylated p38, the active form, increased in the cytoplasm with differentiation (Figure 6A and Additional file 1 Figure S4), suggesting that p38 α might play an important role there. Validation of the phospho-p38 antibody is shown in Additional file 1 Figure S5. Further fractionation of the cytoplasm revealed that p38 α and phospho-p38 were present only in the cytosolic fraction (Figure 6B), meaning that all four of the *in vivo* cytoplasmic phosphorylation sites we identified would be accessible to p38 α . Using Scansite [28], we found that five of the six substrates contained one or more predicted D domains, MAPK docking domains found on many *in vivo* substrates (D domain sites and scores are listed in Additional file 1 Table S4). We next used Genemania [29] to identify previously known associations between these substrates and the p38 activation pathway we recently reviewed and outlined [30]. Four of the substrates have previously known associations with this pathway (Figure



6C), providing additional support that these targets are relevant and that their phosphorylation could be critical for pathway function.

Discussion

We have described here a simple technique for kinase substrate finding that uses whole-cell lysate, can identify sites of phosphorylation in addition to the protein phosphorylated, has the ability to identify both *in vivo* and *in vitro* substrates, and can be used to compare the *in vitro* substrate specificity of two kinases in the same

experiment. Performing the assay with p38 α and p38 β revealed that, although differences in preference are apparent, there are no obvious phosphorylation sites specific to p38 α . However, a major characteristic that distinguishes p38 α from p38 β during myoblast differentiation is localisation, with p38 α located ubiquitously throughout the cell while p38 β is found only at the cell periphery. We also have shown that p38 α can phosphorylate several cytosolic phosphorylation sites *in vivo* and that this may be a critical but previously unknown function for it. These results demonstrate the utility of the substrate screening technique we have developed and how its use not only can find proteins phosphorylated by a kinase of interest but also can answer particular questions and propose novel avenues of research.

The technique we have developed has several advantages over existing substrate finding approaches. It can provide direct substrate identifications and information on *in vivo* relevance in a single experiment. Whole-cell lysate can be used as a source of candidate substrates, allowing screens to be performed on samples of specific relevance to the area of interest instead of on protein or peptide arrays. Our approach for eliminating endogenous kinase activity in lysate is nondenaturing, decreasing the likelihood of false-positive and false-negative substrate phosphorylations that would result from the heat-inactivating approach that others have employed [31,32]. At the same time, the recombinant kinase used in the assay does not require mutation or unnatural ATP analogues as other approaches do [33], making the technique applicable to any kinase that can be made and purified in an active form. Although we used dimethyl labelling for quantitative MS, the technique is fully compatible with iTRAQ, which allows up to eight samples to be compared in a single experiment, meaning that simultaneous *in vivo* substrate identification and kinase comparisons can be performed. Radioactivity and gel work are not required, but can be used to visualise phosphorylations and make qualitative comparisons between kinases on one-dimensional gels. The lysate requirements are also relatively low; 1.5 to 3 mg were used in these assays, which may make the technique difficult to employ with some primary cells lines, but it is easily applicable to most secondary cell lines and to tissues as well. We have treated several different lysate types with FSBA, and complete inhibition of endogenous kinase activity occurred in all, meaning that our approach is likely universally applicable. In this study our approach for identifying *in vivo* substrates utilised an inhibitor, but it is more versatile than that, and can be applied to study phosphorylations triggered by a specific stimulus (using stimulated versus unstimulated cells for lysate pools) without requiring a specific inhibitor be available. Finally, although we have performed

the *in vivo* approach on unfractionated lysate, when combined with fractionation prior to LC-MS/MS, the number of substrates identified could be much greater than what we have demonstrated in principle here.

As mentioned, a major advantage of the technique presented here is that it can be used to compare the substrate profiles of two or more kinases, and we used this to study the substrate specificity of the p38 α and β isoforms. However, we were unable to identify any phosphorylations that were specific to p38 α . There are apparent differences in preference for substrates between p38 α and p38 β , which could be functionally relevant. Assuming that p38 β had the same ubiquitous localisation pattern as p38 α , a low preference for certain critical myogenic phosphorylations could result in a difference in cell behaviour if p38 β were the only isoform present. However, it seems likely that differentiation would still occur, but possibly at a reduced rate or in a compromised way. This is not the case, as p38 β -containing myoblasts that lack p38 α fail to differentiate at all [7-9,34]. Rather, we believe what truly distinguishes the two isoforms is their localisation. p38 α has a known critical role in the nucleus, and our results suggest that the same may be true for the cytoplasm; therefore, p38 β would be unable to compensate because it is found solely at the periphery of the cell. A similar distinction exists for two other closely related kinases, focal adhesion kinase (FAK) and proline-rich tyrosine kinase 2 (PYK2) [35]. In fibroblasts, FAK localises to focal adhesions, whereas PYK2 has a perinuclear distribution. PYK2, which has a kinase domain 60% identical to that of FAK, can compensate for the loss of FAK provided it has a focal adhesion targeting sequence. Together these results suggest that rather than substrate specificity, it is other characteristics, in these cases localisation, that distinguish closely related kinases. Whether the same holds true for the entire kinase family is an intriguing question.

In addition to p38 α 's known role in regulating gene expression at the onset of myoblast differentiation, our results suggest that it is likely to have a cytosolic role as well, possibly in regulating its own pathway. We have found that p38 α is present in the cytosol, active p38 increases in the cytoplasm with differentiation, p38 α can phosphorylate several cytosolic proteins, three of the cytosolic proteins contain predicted D domains, and connections between these proteins and the p38 α activation pathway have been previously reported. Together these results suggest a previously unrecognised role for p38 α in the cytosol.

Conclusions

While the role of the p38 α MAPK during myoblast differentiation has been studied extensively with regard to

gene regulation, our evidence suggests that this kinase is likely involved in other processes as well. The differentiation of myoblasts into a mature myotube involves extensive morphological and functional changes that p38 α might regulate, both through the initiation of a myogenic gene program and through direct phosphorylation and functional modification of cytosolic proteins. No one has previously been able to identify a p38 α -specific substrate, and our large-scale screen suggests these may exist only within the context of cellular localisation. In addition, the substrate screening technique we have developed should serve as a useful tool for studying the other kinases known to be involved in myogenesis, as well as the hundreds of other protein kinases that exist.

Additional material

Additional file 1: Figures S1 through S5 and Tables S1 through S4.

Acknowledgements

JDRK was supported by a Vanier Canada Graduate Scholarship, a Canadian Institutes of Health Research (CIHR) Canada Graduate Scholarship and a Multiple Sclerosis Society of Canada (MSSC) Studentship. LAM holds the Mach Gaensslen Chair in Cardiac Research and was funded by CIHR. ACG is a Canada Research Chair in Functional Proteomics and the Lea Reichmann Chair and is funded by CIHR grant MOP-84314. DF acknowledges a Canada Research Chair in Proteomics and Systems Biology as well as grants from the Natural Sciences and Engineering Research Council of Canada and the J-Louis Lévesque Foundation. RK holds a University Health Research Chair at the University of Ottawa and was supported by grants from CIHR and the MSSC.

Author details

¹Regenerative Medicine Program, Ottawa Hospital Research Institute, 501 Smyth Road, Ottawa, ON, K1H 8L6, Canada. ²Department of Cellular and Molecular Medicine, University of Ottawa, 451 Smyth Road, Ottawa, ON, K1H 8M5, Canada. ³Ottawa Institute of Systems Biology, University of Ottawa, 451 Smyth Road, Ottawa, ON, K1H 8M5, Canada. ⁴Department of Biochemistry, Microbiology and Immunology, University of Ottawa, 451 Smyth Road, Ottawa, ON, K1H 8M5, Canada. ⁵CAS Key Lab of Separation Sciences for Analytical Chemistry, Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian, 116023, China. ⁶Department of Medicine, University of Ottawa, 451 Smyth Road, Ottawa, ON, K1H 8M5, Canada. ⁷Samuel Lunenfeld Research Institute, Mount Sinai Hospital, 600 University Avenue, Toronto, ON, M5G 1X5, Canada. ⁸Department of Molecular Genetics, University of Toronto, 1 King's College Circle, Toronto, ON, M5S 1A8, Canada. ⁹Current address: Samuel Lunenfeld Research Institute, Mount Sinai Hospital, 600 University Avenue, Toronto, ON, M5G 1X5, Canada. ¹⁰Current address: Department of Cancer Biology, Dana-Farber Cancer Institute, 450 Brookline Avenue, and Department of Genetics, Harvard Medical School, 77 Avenue Louis Pasteur, Boston, MA 02115, USA.

Authors' contributions

JDRK and RK conceived of and designed the project. JDRK conceived of the substrate-finding technique. RT assisted JDRK with the implementation of the substrate-finding technique. RECL performed two-dimensional gel work and assisted JDRK with the development of the radioactive approach. FW assisted with quantitative MS analysis and phosphopeptide validation. AB assisted with cell cultures. All other experiments and analysis were performed by JDRK. JDRK wrote the paper, and RK revised and edited it. HZ supervised FW. LAM supervised RECL and gave advice on project design. DF supervised RT and FW, and DF, TP and ACG provided MS facilities and

expertise. RK supervised JDRK. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 22 August 2011 Accepted: 6 March 2012

Published: 6 March 2012

References

- Berwick DC, Tavaré JM: Identifying protein kinase substrates: hunting for the organ-grinder's monkeys. *Trends Biochem Sci* 2004, **29**:227-232.
- Johnson SA, Hunter T: Kinomics: methods for deciphering the kinome. *Nat Methods* 2005, **2**:17-25.
- Manning BD, Cantley LC: Hitting the target: emerging technologies in the search for kinase substrates. *Sci STKE* 2002, **2002**:pe49.
- Sopko R, Andrews BJ: Linking the kinome and phosphorylome: a comprehensive review of approaches to find kinase targets. *Mol Biosyst* 2008, **4**:920-933.
- Holt LJ, Tuch BB, Villén J, Johnson AD, Gygi SP, Morgan DO: Global analysis of Cdk1 substrate phosphorylation sites provides insights into evolution. *Science* 2009, **325**:1682-1686.
- Lluís F, Perdiguero E, Nebreda AR, Muñoz-Cánoves P: Regulation of skeletal muscle gene expression by p38 MAP kinases. *Trends Cell Biol* 2006, **16**:36-44.
- Li Y, Jiang B, Ensign WY, Vogt PK, Han J: Myogenic differentiation requires signalling through both phosphatidylinositol 3-kinase and p38 MAP kinase. *Cell Signal* 2000, **12**:751-757.
- Perdiguero E, Ruiz-Bonilla V, Gresh L, Hui L, Ballestar E, Sousa-Victor P, Baeza-Raja B, Jardí M, Bosch-Comas A, Esteller M, Caelles C, Serrano AL, Wagner EF, Muñoz-Cánoves P: Genetic analysis of p38 MAP kinases in myogenesis: fundamental role of p38 α in abrogating myoblast proliferation. *EMBO J* 2007, **26**:1245-1256.
- Wang H, Xu Q, Xiao F, Jiang Y, Wu Z: Involvement of the p38 mitogen-activated protein kinase α , β , and γ isoforms in myogenic differentiation. *Mol Biol Cell* 2008, **19**:1519-1528.
- Enslin H, Raingeaud J, Davis RJ: Selective activation of p38 mitogen-activated protein (MAP) kinase isoforms by the MAP kinase kinases MKK3 and MKK6. *J Biol Chem* 1998, **273**:1741-1748.
- Boersema PJ, Raijmakers R, Lemeer S, Mohammed S, Heck AJ: Multiplex peptide stable isotope dimethyl labeling for quantitative proteomics. *Nat Protoc* 2009, **4**:484-494.
- Li QR, Ning ZB, Tang JS, Nie S, Zeng R: Effect of peptide-to-TiO₂ beads ratio on phosphopeptide enrichment selectivity. *J Proteome Res* 2009, **8**:5375-5381.
- Wang F, Dong J, Jiang X, Ye M, Zou H: Capillary trap column with strong cation-exchange monolith for automated shotgun proteome analysis. *Anal Chem* 2007, **79**:6599-6606.
- Wang F, Chen R, Zhu J, Sun D, Song C, Wu Y, Ye M, Wang L, Zou H: A fully automated system with online sample loading, isotope dimethyl labeling and multidimensional separation for high-throughput quantitative proteome analysis. *Anal Chem* 2010, **82**:3007-3015.
- Olsen JV, Blagoev B, Gnäd F, Macek B, Kumar C, Mortensen P, Mann M: Global, *in vivo*, and site-specific phosphorylation dynamics in signaling networks. *Cell* 2006, **127**:635-648.
- Crooks GE, Hon G, Chandonia JM, Brenner SE: WebLogo: a sequence logo generator. *Genome Res* 2004, **14**:1188-1190.
- Hashimoto E, Takio K, Krebs EG: Amino acid sequence at the ATP-binding site of cGMP-dependent protein kinase. *J Biol Chem* 1982, **257**:727-733.
- Kamps MP, Taylor SS, Sefton BM: Direct evidence that oncogenic tyrosine kinases and cyclic AMP-dependent protein kinase have homologous ATP-binding sites. *Nature* 1984, **310**:589-592.
- Russo MW, Lukas TJ, Cohen S, Staros JV: Identification of residues in the nucleotide binding site of the epidermal growth factor receptor/kinase. *J Biol Chem* 1985, **260**:5205-5208.
- Zoller MJ, Nelson NC, Taylor SS: Affinity labeling of cAMP-dependent protein kinase with *p*-fluorosulfonylbenzoyl adenosine: covalent modification of lysine 71. *J Biol Chem* 1981, **256**:10837-10842.
- Knight JD, Qian B, Baker D, Kothary R: Conservation, variability and the modeling of active protein kinases. *PLoS One* 2007, **2**:e982.

22. Schwartz D, Gygi SP: An iterative statistical approach to the identification of protein phosphorylation motifs from large-scale data sets. *Nat Biotechnol* 2005, **23**:1391-1398.
23. Manning G, Whyte DB, Martinez R, Hunter T, Sudarsanam S: The protein kinase complement of the human genome. *Science* 2002, **298**:1912-1934.
24. Andrés V, Walsh K: Myogenin expression, cell cycle withdrawal, and phenotypic differentiation are temporally separable events that precede cell fusion upon myogenesis. *J Cell Biol* 1996, **132**:657-666.
25. Diebold ML, Koch M, Loeliger E, Cura V, Winston F, Cavarelli J, Romier C: The structure of an lws1/Spt6 complex reveals an interaction domain conserved in TFIIIS, Elongin A and Med26. *EMBO J* 2010, **29**:3979-3991.
26. Yoh SM, Cho H, Pickle L, Evans RM, Jones KA: The Spt6 SH2 domain binds Ser2-P RNAPII to direct lws1-dependent mRNA splicing and export. *Genes Dev* 2007, **21**:160-174.
27. Asally M, Yasuda Y, Oka M, Otsuka S, Yoshimura SH, Takeyasu K, Yoneda Y: Nup358, a nucleoporin, functions as a key determinant of the nuclear pore complex structure remodeling during skeletal myogenesis. *FEBS J* 2011, **278**:610-621.
28. Obenauer JC, Cantley LC, Yaffe MB: Scansite 2.0: Proteome-wide prediction of cell signaling interactions using short sequence motifs. *Nucleic Acids Res* 2003, **31**:3635-3641.
29. Warde-Farley D, Donaldson SL, Comes O, Zuberi K, Badrawi R, Chao P, Franz M, Grouios C, Kazi F, Lopes CT, Maitland A, Mostafavi S, Montojo J, Shao Q, Wright G, Bader GD, Morris Q: The GeneMANIA prediction server: biological network integration for gene prioritization and predicting gene function. *Nucleic Acids Res* 2010, **38** Web server: W214-W220.
30. Knight JD, Kothary R: The myogenic kinase: protein kinases critical to mammalian skeletal myogenesis. *Skelet Muscle* 2011, **1**:29.
31. Huang SY, Tsai ML, Chen GY, Wu CJ, Chen SH: A systematic MS-based approach for identifying *in vitro* substrates of PKA and PKG in rat uteri. *J Proteome Res* 2007, **6**:2674-2684.
32. Troiani S, Uggeri M, Moll J, Isacchi A, Kalisz HM, Rusconi L, Valsasina B: Searching for biomarkers of Aurora-A kinase activity: identification of *in vitro* substrates through a modified KESTREL approach. *J Proteome Res* 2005, **4**:1296-1303.
33. Blethrow JD, Glavy JS, Morgan DO, Shokat KM: Covalent capture of kinase-specific phosphopeptides reveals Cdk1-cyclin B substrates. *Proc Natl Acad Sci USA* 2008, **105**:1442-1447.
34. Ruiz-Bonilla V, Perdiguero E, Gresh L, Serrano AL, Zamora M, Sousa-Victor P, Jardí M, Wagner EF, Muñoz-Cánoves P: Efficient adult skeletal muscle regeneration in mice deficient in p38 β , p38 γ and p38 δ MAP kinases. *Cell Cycle* 2008, **7**:2208-2214.
35. Klingbeil CK, Hauck CR, Hsia DA, Jones KC, Reider SR, Schlaepfer DD: Targeting Pyk2 to β 1-integrin-containing focal contacts rescues fibronectin-stimulated signaling and haptotactic motility defects of focal adhesion kinase-null cells. *J Cell Biol* 2001, **152**:97-110.

doi:10.1186/2044-5040-2-5

Cite this article as: Knight *et al.*: A novel whole-cell lysate kinase assay identifies substrates of the p38 MAPK in differentiating myoblasts. *Skeletal Muscle* 2012 **2**:5.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

